

Injection Tips

for Buprenorphine ER and Meloxicam ER

Important notes for administration and storage:

- ZooPharm's ER formulations are viscous and should at least be brought to room temperature for 30-60 minutes prior to use.
- Do NOT store ER formulations in filled syringes of any type, even if under refrigeration. Interaction with syringe components could cause a chemical reaction that both changes viscosity and properties of both Buprenorphine ER and Meloxicam ER. (e.g. polymerization that turns the formulation cloudy, whitish and even possibly to solidify).
- Some clinics have reported that placing the Buprenorphine ER vial in a warm water bath and bringing it within one degree of animal body temperature prior to injection, reduces viscosity, makes it easier to draw-up and slowly inject... decreasing animal discomfort during administration.

Please consult with a veterinarian for proper use.

**It is important to use polypropylene syringes,
not polycarbonate, with all ER formulations.**

(see reverse side for examples)



Examples of Polypropylene Syringes

- Sterile 1ml syringe
- Clear Polypropylene (PP) barrel
- Non-toxic, Non-pyrogenic and latex free
- Reduced dead space for reduced waste
- Smooth plunger action for greater filling accuracy
- Graduated scale in 0.01ml increments



1cc TUBERCULIN SYRINGE ONLY

Catalog No.	Description
26048*	1cc Slip Tip with Cap
26049*	1cc Lock Tip with Cap
26050	1cc Lock Tip

Recommended



We recommend EXELint Model # 26049.

Visit bit.ly/zp-syringe to order!

Examples of Polycarbonate Syringes (Do NOT use)



BD Luer-Lock 1ml Syringe (Catalog No. 309628)

- Crystal-clear polycarbonate barrel
- integral Luer-Lock hub
- Readily visible fluid level

ZooPharm does not make claims for the safety or efficacy of our compounded medications. © ZooPharm 2022. All rights reserved. 1022